ON THE GENERATION OF STRONG MAGNETIC FIELDS IN THE INTERACTION OF ULTRAINTENSE SHORT LASER PULSE WITH AN OVERDENSE PLASMA TARGET

V.I. Karas^a, O.V. Batishchev^b, M. Bornatici^c

^aNSC "Kharkov Institute of Physics & Technology", Kharkov 61108, Ukraine ^bMIT Plasma Fusion Center, Cambridge, Massachusetts 02139, USA ^cINFM, Dipartimento di Fisica "A. Volta", Università degli Studi di Pavia, Pavia 27100, Italy

The interaction between intense laser radiation and matter is known to produce a wealth of nonlinear effects. Those include fast electron and ion generation [1-5] indicating that ultra-strong electric fields are produced in the course of the laser – plasma interaction. An equally ubiquitous, although less studied, effect accompanying laser – matter interaction is the generation of ultra-strong magnetic fields in the plasma [6–11]. Magnetic fields can have a significant effect on the overall nonlinear plasma dynamics. Extremely high (few megagauss) azimuthal magnetic fields play an essential role in the particle transport, propagation of laser pulses, laser beam self - focusing, penetration of laser radiation into the overdense plasma and the plasma electron and ion acceleration.

By means of a 2.5 - dimensional numerical simulation on the macroparticles method it is possible to find the magnetic field spatial and temporal distribution without usage an adapted parameter unlike of the conventional $\nabla_{nx}\nabla T$ mechanism (see for example [6, 32]). On the other hand, theoretical model for the generation of a magnetic field proposed by Sudan [8] is not appropriate, this model being very large ratio of plasma density to critical density and when the $\nabla_{nx}\nabla T$ contribution is not relevant.

^aNSC "Kharkov Institute of Physics & Technology", Kharkov 61108, Ukraine ^bMIT Plasma Fusion Center, Cambridge, Massachusetts 02139, USA ^cINFM, Dipartimento di Fisica "A. Volta", Università degli Studi di Pavia,Pavia 27100, Italy

1. The generation of an axial magnetic field in the plasma by a circularly (or elliptically) polarized laser is often referred to as the inverse faraday effect (ife). First theoretically described by Pitaevskii [12] and Steiger and Woods [13] it results from the features of the electron motion in a circularly polarized electromagnetic wave. During the interaction of the plasma electrons with the circularly polarized laser pulse, electrons absorb both the laser energy and the angular momentum of the laser pulse. In particular,

THE ANGULAR MOMENTUM ABSORPTION LEADS TO THE ELECTRON ROTATION AND GENERATION OF THE AXIAL MAGNETIC FIELD BY THE AZIMUTHAL ELECTRON CURRENT.

Naturally, IFE does not occur for a linearly polarized laser pulse since it does not possess any angular momentum. IFE has since been measured in several experiments [9,14, 15]. The conditions under which ife is possible are still not fully explored. What is theoretically known [16] is that THERE IS NO MAGNETIC-FIELD GENERATION DURING THE INTERACTION OF THE INHOMOGENEOUS CIRCULARLY POLARIZED ELECTROMAGNETIC WAVES WITH THE HOMOGENEOUS PLASMA.

Magnetic field can be produced in the presence the strong plasma inhomogeneity [17–19] either preformed or developed self-consistently during the interaction. More recently a mechanism has been proposed for the generation of an axial magnetic field through the transfer of the spin of the photons during the absorption of a transversely nonuniform circularly polarized radiation [20].

THE MAGNETIC FIELD THUS GENERATED HAS A MAGNI-TUDE PROPORTIONAL TO THE TRANSVERSE GRADIENT OF THE ABSORBED INTENSITY AND INVERSE PROPORTIO-NAL TO THE ELECTRON DENSITY, THE LATTER SEALING BEING IN CONTRAST TO EARLIER THEORIES OF IFE [20].

Along the same lines as [20], it has been demonstrated that in laserplasma interactions strong axial magnetic fields can be generated through angular (spin) momentum absorption by either electron-ion collisions or ionization.

^aNSC "Kharkov Institute of Physics & Technology", Kharkov 61108, Ukraine

- ^bMIT Plasma Fusion Center, Cambridge, Massachusetts 02139, USA
- ^cINFM, Dipartimento di Fisica "A. Volta", Università degli Studi di Pavia, Pavia 27100, Italy

Workshop "Ultrashort pulse laser-plasma interactions" and INTAS-01-233 project meeting, (SEPTEMBER 23-27, 2003)

Yet another mechanism of magnetic field generation has been proposed (in framework of classical electrodynamics) [21], based on the resonant absorption by energetic electrons of a circularly polarized laser pulse. The resonance occurs between the fast electrons, executing transverse (betatron) oscillations in a fully or partially evacuated plasma channel, and the electric field of the laser pulse. The betatron oscillations are caused by the action of the electrostatic force of the channel ions and selfgenerated magnetic field. This type of resonant interaction was recently suggested as a mechanism for accelerating electrons to highly relativistic energies [21,22].

WHEN A CIRCULARLY POLARIZED LASER PULSE IS EM-PLOYED, ITS ANGULAR MOMENTUM CAN BE TRANSFER-RED TO FAST RESONANT ELECTRONS ALONG WITH ITS ENERGY. THE RESULTING ELECTRON BEAM SPIRALS AROUND THE DIRECTION OF THE LASER PROPAGATION, GENERATING THE AXIAL MAGNETIC FIELD [21].

In [21] the intensity of the magnetic field generated in relativistic laser channel was calculated taking into account self-generated static fields, which are neglected in known IFE theories [17–19]. Calculations [21] are in agreement with the recent experiments at the Rutherford Appleton Laboratory (RAL) [15] which exhibited very large (several megagauss) axial magnetic fields during the propagation of a sub-picosecond laser pulse in a tenuous plasma. The relevant aspect of the RAL experiment is that both fast electrons and the strong magnetic field were measured in the same experiment.

2. In an underdense longitudinal inhomogeneous plasma the nonrelativistic two-dimensional treatment of self-generated magnetic fields was presented in the article [23], showing that a laser beam propagating along a density plasma gradient produces a rotational current which gives rise to a quasi-static magnetic field. An analogous mechanism was considered earlier in the paper [24] whereas the nonlinear mixing of two electromagnetic waves in a nonuniform plasma was discussed in the work [25]. They investigated a circularly polarized pulse for which the generation of low-frequency electromagnetic field is due to the inverse faraday effect. In the extremely strong relativistic regime the magnetic

^aNSC "Kharkov Institute of Physics & Technology", Kharkov 61108, Ukraine

^bMIT Plasma Fusion Center, Cambridge, Massachusetts 02139, USA

^cINFM, Dipartimento di Fisica "A. Volta", Università degli Studi di Pavia,Pavia 27100, Italy

field generated by the laser beams in underdense plasma was recently studied numerically [26]. The main objective of the work [18] was to investigate self-generated quasi-static magnetic fields both in the laser pulse body and behind the pulse in the region of the wake-field. Authors treated the laser radiation as linearly polarized and the plasma as uniform and underdense. The analytical work was based on a perturbation theory applied to the set of relativistic cold electron fluid equations and Maxwell's equations. The quasi-static magnetic field generated by a short laser pulse in a uniform rarefied plasma is found analytically and compared to two-dimensional particle-in-cell simulations.

IT IS SHOWN THAT A SELF-GENERATION OF QUASISTA-TIC MAGNETIC FIELDS TAKES PLACE IN FOURTH ORDER WITH RESPECT TO THE PARAMETER V_E/C WHERE V_E AND C ARE THE ELECTRON QUIVER VELOCITY AND THE LIGHT SPEED, RESPECTIVELY. IN THE WAKE REGION THE MAGNETIC FIELD POSSESSES A COMPONENT WHICH IS HOMOGENEOUS IN THE LONGITUDINAL DIRECTION AND IS DUE TO THE STEADY CURRENT PRODUCED BY THE PLASMA WAKEFIELD AND A COMPONENT WHICH IS OSCILLATING AT THE WAVE NUMBER $2K_B$

Where k_p is the wave number of the plasma wake, a known property of nonlinear plasma waves [27,28]. Numerical particle simulations confirm the analytical results and are also used to treat the case of high intense laser pulses with $v_e/c > 1$. The resultant magnetic field has a focusing effect on relativistic electrons in the plasma wakefield accelerator context.

3. Further we discuss the physical mechanism for generation of very high "quasi-static" magnetic fields in the interaction of an ultraintense short laser pulse with an overdense plasma target owing to the spatial gradients and non-stationary character of the ponderomotive force. Numerical (particle-in-cell) simulations by Wilks et al. [10] of the INTERACTION OF AN ULTRAINTENSE LASER PULSE WITH AN OVERDENSE PLASMA TARGET HAVE REVEALED NON-OSCILLATORY SELF-GENERATED MAGNETIC FIELDS UP TO 250 MG IN THE OVERDENSE PLASMA, THAT THIS

^aNSC "Kharkov Institute of Physics & Technology", Kharkov 61108, Ukraine ^bMIT Plasma Fusion Center, Cambridge, Massachusetts 02139, USA ^cINFM, Dipartimento di Fisica "A. Volta", Università degli Studi di Pavia,Pavia 27100, Italy

NONOSCILLATORY MAGNETIC FIELD AROUND THE HEATED SPOT IN THE CENTER OF THE PLASMA, THE MAGNETIC FIELD GENERATION BEING ATTRIBUTED TO THE ELECTRON HEATING AT THE RADIATION-PLASMA INTERFACE.

The spatial and temporal evolution of spontaneous megagauss magnetic fields, generated during the interaction of a picosecond pulse with solid targets at irradiances above 5×10^{18} W/cm² have been measured using Faraday rotation with picosecond resolution, the observations being limited to the region of underdense plasma and after a laser pulse fig. 1, [6]. A high density plasma jet has been observed simultaneously with the magnetic fields by interferometry and optical emission.

Because of the high temporal resolution of the probe diagnostic, a quantitative measurement of the transient nature of the fields has been obtained for the first time. Interestingly, no Faraday rotation was detected immediately after the interaction, a possible reason being that the fields were still limited to regions not accessible for probing. After 5 ps the typical signatures corresponding to a toroidal field surrounding the laser axis (i.e., a dark and a bright pattern on opposing sides of the axis, in the proximity of the target surface) began to appear.

The sense of rotation is the same as observed in previous measurements in longer pulse regimes and is consistent with fields generated by the thermoelectric mechanism (see for example [30]).

In paper [7] the first direct measurements of high energy proton generation (up to **18 MeV**) and propagation into a solid target during such intense laser plasma interactions were reported. Measurements of the deflection of these energetic protons were carried out which imply that magnetic fields in excess of **30 MG** exist inside the target. The structure of these fields is consistent with those produced by a beam of hot electrons which has also been measured in these experiments. Such observations are the first evidence of the large magnetic fields which have been predicted to occur during laser-target interactions in dense plasma [10]. The intensity on target of laser radiation was up to **5 x 10¹⁹** W/cm² and was determined by simultaneous measurements of the laser pulse energy, duration, and focal spot size.

^aNSC "Kharkov Institute of Physics & Technology", Kharkov 61108, Ukraine

^bMIT Plasma Fusion Center, Cambridge, Massachusetts 02139, USA

^cINFM, Dipartimento di Fisica "A. Volta", Università degli Studi di Pavia,Pavia 27100, Italy

IN [29] THE PROBLEM OF HIGH-INTENSITY, LINEARLY POLARIZED ELECTROMAGNETIC PULSE INCIDENT ONTO A COLLISIONLESS PLASMA LAYER IS SOLVED NUMERICALLY CARTESIAN COORDINATE SYSTEM IN A IN A 2.5-D FORMULATION (Z IS THE CYCLIC COORDINATE AND THERE ARE THREE COMPONENTS OF THE MOMENTUM) BY MEANS COMPASS (COMPUTER PLASMA AND SURFACE SIMULATION) CODE. THE RECENT REVIEW [30] AND REFERENCES THEREIN DETAILED INFORMATION COMBINE Α CONCERNING COMPASS CODE AS WELL AS ITS POSSIBILITIES AND APPLICATIONS. ONE OF GENERAL ADVANTAGE OF THE COMPLETE NUMERICAL SIMULATION IS POSSIBILITY AS WELL AS AT THE LABORATORY EXPERIMENTS OBTAIN ALL NECESSARY INFORMATION CONCERNING SPATIAL AND TEMPORAL DYNAMICS OF BOTH PARTICLES AND SELF-CONSISTENT ELECTROMAGNETIC FIELDS WITHOUT USAGE OF ADDITIONAL DATA (REFLECTION AND ABSORPTION CHANGES OF COEFFICIENTS, EITHER Α PLASMA TEMPERATURE OR DIFFERENT PLASMA PARAMETERS) IN SITUATION AT INTERACTION **INTENSIVE** GIVEN Α ELECTROMAGNETIC PULSE WITH PLASMAS. WE GIVE ONLY THE EXTERNAL PARAMETERS, THE BOTH INITIAL AND BOUNDED CONDITIONS FOR PARTICLES AND FIELDS AND AS RESULTS OF A NUMERICAL SIMULATION WE ATTAIN ALL CHARACTERICTICS OF PLASMAS TOGETHER WITH PULSED SELF-CONSISTENT ELECTROMAGNETIC FIELDS. IN GIVEN PROBLEM WE CONSIDER AT THE INITIAL TIME, A COLD **TWO-COMPONENT** NEUTRAL (IONS MOTIONLESS AND ELECTRONS WITH A REAL RATIO OF THEIR MASSES) PLASMA WITH UNIFORM DENSITY FILLS THE WHOLE RIGHT-HAND PART OF RECTANGULAR DOMAIN X x Y = L_x x L_y = 128 x 64 c/w_{pe} (C IS THE SPEED OF LIGHT, w_{pe} IS THE ELECTRON FREQUENCY). THE LINE X 40 PLASMA L_B == C/W_{pe} PLASMA-VACUUM BOUNDARY. REPRESENTS THE THE

^aNSC "Kharkov Institute of Physics & Technology", Kharkov 61108, Ukraine

^bMIT Plasma Fusion Center, Cambridge, Massachusetts 02139, USA

^cINFM, Dipartimento di Fisica "A. Volta", Università degli Studi di Pavia,Pavia 27100, Italy

SYSTEM IS PERIODICAL ALONG THE Y-COORDINATE; THE PLASMA PARTICLES ARE REFLECTED ELASTICALLY FROM THE RIGHT-HAND AND LEFT-HAND BOUNDARIES. THE BOUNDARY CONDITIONS INITIAL AND FOR THE ELECTROMAGNETIC FIELDS (THEY ARE MEASURED IN UNITS $m_{e}w_{0}c/e$) ARE THE FOLLOWING: E (T<0) = B (T<0) = 0; E_Y (X=0, Y, T>0 = B_Z (X=0, Y, T>0) = A (Y)COS (w_0 T); A (Y) = A_0EXP ((Y- $L_{\rm Y}/L_0)^2$, WHERE $w_0 = w_{pe}/2$ IS THE FREQUENCY OF THE INCIDENT ELECTROMAGNETIC RADIATION. THE RADIATION INTENSITY AD PROVIDES A KINETIC MOMENTUM OF ABOUT $3M_{\rm E}C$ TO BE CARRIED BY OSCILLATIONS; $L_0 = 10 c/w_{\rm ne}$.

THE MOST CHARACTERISTIC FEATURE OF THE ACTION OF AN INTENSE, NORMALLY INCIDENT ELECTROMAGNETIC PULSE ONTO AN ULTRAHIGH-DENSITY PLASMA CONSISTS IN A "WELL-DIGGING" EFFECT. THE DEPTH OF THE "WELL" IN THE PLASMA PROFILE INCREASES WITH TIME AND IS 15 c/w_{pe} AT TIME 500 w_{pe}^{-1} . WORTH NOTHING IS THE GROWING IN TIME SHARP NONUNIFORMITY OF THE PERTURBED PLASMA LAYER IN THE TRANSVERSE DIRECTION.

MAGNETIC SHOWN THAT THE THEREBY IT FIELD OSCILLATES WITH THE DOUBLED FREQUENCY OF A LASER RADIATIAN, BUT IT HAVE UNCHANGED DIRECTION. THE MAGNETIC FIELD OSCILLATIONS ARE SAW GOOD AT FIG. 2. NOW LET US LOOK AT FIGS. 2a AND 2b, WHICH PRESENT TWO INSTANTANEOUS MAGNETIC FIELD B_Z DISTRIBUTIONS SEPARATED IN THE TIME BY pw_{pe}^{-1} (A PLASMA WAVE HALF-IT WILL BE NOTED THAT A MAXIMUM OF PERIOD). MAGNETIC FIELD (1.1) IN THE POINT (38,31) IN THE TIME tw_{pe} =200 (SEE FIG.2A) AFTER A PLASMA WAVE HALF-PERIOD (SEE FIG. 2B) REPLACED IN THIS POINT VERY LOW VALUE OF MAGNETIC FIELD.

^aNSC "Kharkov Institute of Physics & Technology", Kharkov 61108, Ukraine ^bMIT Plasma Fusion Center, Cambridge, Massachusetts 02139, USA ^cINFM, Dipartimento di Fisica "A. Volta", Università degli Studi di Pavia,Pavia 27100, Italy THE MAGNETIC FIELD B_z ATTAINS ITS PEAK VALUE OF (0.64) AT THE POINT (36.5, 29), CF. FIG. 2b. IT IS SHOWN THAT THE MAGNETIC FIELD BZ IS CONSISTENT WITH A LINEARLY MODULATED CURRENT FLOWING ALONG THE LINE $Y = L_{x}/2$. DO NOT OBSERVE THIS FIELD TO CHANGE ITS WE DIRECTION. BUT ITS STRENGTH VARIES SIGNIFICANTLY IN HENCE. THE MAGNETIC TIME. FIELD CANNOT BE CONSIDERED AS OUASI-STATIC BECAUSE IT VARIES BY MORE THAN AN ORDER OF MAGNITUDE OVER A TIME OF $2pw_{pe}^{-1}$.

THE MAGNITUDE OF THE "DC" MAGNETIC FIELD IS TEN TIMES AS LOW AS THE MAXIMUM MAGNETIC FIELD. ONE SHOULD NOTE THAT THE NUMERICAL SIMULATION HAS BEEN MADE UNDER VERY OPTIMAL CONDITIONS: UNIFORM PLASMA DENSITY MAKES IT SURE A OWN PLASMA LONGITUDINAL OSCILLATION RESONANCE WITH A MODULATION DENSITY OF PARTICLES IN A WAVE AS WELL AS A MAXIMUM FREQUENCY OF NONLINEAR TOMSON SCATTERING SPECTRUM. IN EXPERIMENTS A PLASMA INHOMOGENEITY WAS VERY ESSENTIAL. WITH THE RESULT THAT RESONANT CONDITIONS WERE FULFILLED ONLY IN A SMALL PLASMA REGION. THE RESULTS OF OUR NUMERICAL SIMULATION WHEN RATIO OF PLASMA DENSITY TO CRITICAL DENSITY IS EQUAL 9 (IN THIS CASE THE **RESONANT CONDITIONS ARE NOT SATISFIED (SEE FIG. 3)) TO** BRING IN EVIDENCE OF A RESONANCE SIGNIFICANCE. AFTERWARDS THE INTERACTION PULSE, ONLY THE "DC" EXISTS, MAGNETIC FIELD AS MEASURED IN THE UNDERDENSE PLASMA REGION IN [6]. ON THE BASIS OF THE FORMULA

 $B_{dc}(MG) = 4.2x(10^{-22}I(W/m^2))^{1/2}(I(mn))^{-1}$

(WHERE I IS THE INTENSITY OF THE INCIDENT LASER RADIATION) ONE OBTAINS A "DC" MAGNETIC FIELD

^cINFM, Dipartimento di Fisica "A. Volta", Università degli Studi di Pavia, Pavia 27100, Italy

^aNSC "Kharkov Institute of Physics & Technology", Kharkov 61108, Ukraine

^bMIT Plasma Fusion Center, Cambridge, Massachusetts 02139, USA

MAGNITUDE OF FEW MG FOR THE EXPERIMENTAL PARAMETERS OF [6], AND A FEW TENS MG FOR THE EXPERIMENTAL CONDITIONS OF [7]. A DIFFERENCE STILL ON ORDER OF VALUE IS CONDITIONED THAT AT SUCH INTENSITIES ONLY 10% OF THE INCIDENT LASER RADIATION IS ABSORBED IN AGREEMENT WITH [31].

MEANS OF A 2.5 - DIMENSIONAL BY NUMERICAL SIMULATION ON THE MACROPARTICLES METHOD IT IS POSSIBLE TO FIND THE MAGNETIC FIELD SPATIAL AND TEMPORAL DISTRIBUTION WITHOUT USAGE AN ADAPTED PARAMETER UNLIKE OF THE CONVENTIONAL $\nabla n x \nabla T$ MECHANISM (SEE FOR EXAMPLE [6, 32]). ON THE OTHER HAND, THEORETICAL MODEL FOR THE GENERATION OF A MAGNETIC FIELD PROPOSED BY SUDAN [8] IS NOT APPROPRIATE, THIS MODEL BEING VERY LARGE RATIO OF PLASMA DENSITY TO CRITICAL DENSITY AND WHEN THE $\nabla nx \nabla T$ CONTRIBUTION IS NOT RELEVANT.

THE WORK BY V.I. KARAS`WAS SUPPORTED IN PART BY THE CARIPLO FOUNDATION (COMO, ITALY) AND INTAS PROJECT 01-233.

REFERENCES

 A. MODENA, A. E. DANGOR, Z. NAJMUDIN, C. E. CLAYTON, K. MARSH, C. JOSHI, V. MALKA, C.B. DARROW, C. DANSON, D. NEELY, AND F. N. WALSH, NATURE (LONDON) **377**, 606 (1995).
A. TING, C. I. MOORE, K. KRUSHELNICK, C. MANKA, E. ESAREY, P. SPRANGLE, R. HUBBARD, H. R. BURRIS, R. FISCHER, AND M. BAINE, PHYS. PLASMAS **4**, 1889 (1997).
R. WAGNER, S.-Y. CHEN, A. MAKSIMCHUK, AND D. UMSTADTER, PHYS. REV. LETT. **78**, 3125 (1997).
K. KRUSHELNICK, E. L. CLARK, Z. NAJMUDIN, M. SALVATI, M. I. K. SANTALA, M. TATARAKIS, A. E. DANGOR, V. MALKA, D. NEELY, R. ALLOTT, AND C. DANSON, PHYS. REV. LETT. **83**, 737 (1999).

^aNSC "Kharkov Institute of Physics & Technology", Kharkov 61108, Ukraine ^bMIT Plasma Fusion Center, Cambridge, Massachusetts 02139, USA ^cINFM, Dipartimento di Fisica "A. Volta", Università degli Studi di Pavia,Pavia 27100, Italy [5] A. MAKSIMCHUK, S. GU, K. FLIPPO, D. UMSTADTER, AND

V. YU. BYCHENKOV, PHYS. REV. LETT. 84, 4180 (2000).

[6] M. BORGHESI, A. J. MACKINNON, A. R. BELL, R. GAILARD, AND O. WILLI, PHYS.REV. LETT. **81**,112 (1998).

[7] E. L. CLARK, K. KRUSHELNICK , J. R. DAVIES, M. ZEPF, M. TATARAKIS, F. N. BEG, A. MACHCEK, P. A. NORREYS, M. I. K. SANTALA, I. WATTS, AND A. E. DANGOR, PHYS. REV. LETT. **84**, 670 (2000).

[8] R. N. SUDAN, PHYS. REV. LETT. 20, 3075 (1993).

[9] A. PUKHOV, AND J. MEYER-TER-VEHN, PHYS. REV. LETT. 76, 3975 (1999); A. PUKHOV, AND J. MEYER-TER-VEHN, PHYS. PLASMAS 5, 1880 (1998); Y. HOROVITZ, S. ELIEZER, A. LUDMIRSKY, Z. HENIS, E. MOSHE, R. SPITALNIK, AND B. ARAD, PHYS. REV. LETT. 78, 1707 (1997).

[10] S. C. WILKS, W. L. KRUER, M. TABAK, AND A. B. LANGDON, PHYS. REV. LETT. **69**, 1383 (1992); S.G. WILKS, PHYS. FLUIDS **B5**, 2603 (1993); R. J. MASON AND M. TABAK, IBID. **80**, 524 (1998).

[11] L. M. GORBUNOV, P. MORA, AND T. M. ANTONSEN, PHYS. PLASMAS 4, 4358 (1997).

[12] L. P. PITAEVSKII, SOV. PHYS. JETP 12, 1008 (1961).

[13] J. DESCHAMOS ET AL., PHYS. REV. LETT. 25, 1330 (1970); A.D. STEIGER AND C. H. WOODS, PHYS. REV. A 5, 1467(1972).

[14] T. LEHNER, EUROPHYS. LETT. **50**, 480 (2000).

[15] Z. NAJMUDIN, M. TATARAKIS, E. PUKHOV, E. L. CLARK, R. J. CLARK, A. E. DANGOR, J. FAURE V. MALKA, D. NEELY, M. I. K. SANTALA, AND K. KRUSHELNICK, PHYS. REV. LETT. **87**, 215004 (2001).

[16] A. SH. ABDULLAEV AND A. A. FROLOV, SOV. PHYS. JETP **54**, 493 (1981).

[17] Z. H. SHENG AND J. MEYER-TER-VEHN, PHYS. REV. E **54**, 1833 (1996).

[18] L. M. GORBUNOV, P. MORA, AND T. M. ANTONSEN, JR., PHYS. REV. LETT. **76**, 2495 (1996).

^aNSC "Kharkov Institute of Physics & Technology", Kharkov 61108, Ukraine

^bMIT Plasma Fusion Center, Cambridge, Massachusetts 02139, USA

^cINFM, Dipartimento di Fisica "A. Volta", Università degli Studi di Pavia, Pavia 27100, Italy

Workshop "Ultrashort pulse laser-plasma interactions" and INTAS-01-233 project meeting, (SEPTEMBER 23-27, 2003)

[19] V. I. BEREZHIANI, S. M. MAHAJAN, AND N. L.

SHATASHVILI, PHYS. REV. E 55, 995 (1997).

[20] M. G. HAINES, PHYS. REV. LETT. 87, 135005 (2001).

[21] I.YU. KOSTYUKOV, G. SHVETS, N. J. FISCH, AND J. M. RAX,

LASER PART. BEAMS 19, 133 (2001); BULL. AM. PHYS. SOC. 47,

39 (2000);I.YU. KOSTYUKOV, G. SHVETS, N. J. FISCH, AND J. M.

RAX, PHYS. PLASMAS **9**, 636 (2002); G. SHVETS, N.J. FISCH, AND J.-M. RAX, PHYS. REV. E**65**, 046403 (2002).

[22] A. PUKHOV, Z. M. SHENG, AND J. MEYER-TER-VEHN,

PHYS. PLASMAS 6, 2847 (1999); C. GAHN, G. TSAKIRIS, A.

PUKHOV ET AL., PHYS. REV. LETT. 83, 4772 (1999).

[23] V. K. TRIPATHI AND C. S. LIU, PHYS. PLASMAS **1**, 990 (1994).

[24] K. B. DYSTHE, E. MJOLHUS, AND J. TRULSEN, J. GEOPHYS. RES. **83**, 1985 (1978).

[25] V. YU BYCHENKOV, V. I. DEMIN, AND V. T. TIKHONCHUK, SOV. PHYS. JETP **78**, 62 (1994).

[26] G. A. ASKAR'YAN, S. V. BULANOV, F. PEGORARO, AND A. M. PUKHOV, PIS'MA ZH. EKSP. TEOR. FIZ. **60**, 240 (1994) [JETP LETT. **60**, 251 (1994)].

[27] A. R. BELL AND P. GIBBON, PLASMA PHYS. CONTROLLED FUSION **30**, 1319 (1988).

[28] G. MIANO, PHYS. SCR. T**30**, 198 (1990).

[29] O. V. BATISHCHEV, V. I. KARAS', V. D. LEVCHENKO, AND YU. S. SIGOV, PLASMA PHYS. REPORTS **20,** 587 (1994).

[30] V.A. BALAKIREV V.I. KARAS', AND I.V. KARAS', PLASMA PHYSICS REPORTS **28,** 125 (2002).

[31] D. F. PRICE, R. M. MORE, R. S. WALLING, G. GUETHLEIN, R. L. SHEPHERD, R.E. STEWART, AND W. E. WHITE, PHYS. REV. LETT. **75**, 252 (1995).

[32] M. G. HAINES, PHYS. REV. LETT. 78, 254 (1997).

^aNSC "Kharkov Institute of Physics & Technology", Kharkov 61108, Ukraine

^bMIT Plasma Fusion Center, Cambridge, Massachusetts 02139, USA

^cINFM, Dipartimento di Fisica "A. Volta", Università degli Studi di Pavia,Pavia 27100, Italy